

Antibiotic resistance associated with aquaculture in Vietnam



Hoang Nam Kha Nguyen^A, Thi Thu Hao Van^{B,C} and Peter J Coloe^B

^AFaculty of Fisheries, Nong Lam University, Thu Duc District, Ho Chi Minh City, Vietnam

^BSchool of Science, RMIT University, Bundoora West Campus, Bundoora, Vic. 3083, Australia

^CEmail: thithuhao.van@rmit.edu.au

The fishery sector is an important source of income, employment and food supply in Vietnam. In 2014, Vietnam was ranked the world's fourth largest exporter and the third largest producer of farmed food fish. Vietnam seafood export has attained the value of over US\$6.0 billion since 2011 and reached a peak of US\$7.9 billion in 2014. However, many problems and diseases confront sustainable development of the fishery sector and overuse of antibiotics is considered a major challenge. Antibiotics are used in aquaculture for both therapeutic and prophylactic reasons. Various antimicrobials used in human medicine are also used for food animals even for non-therapeutic use. The use of antibiotics in health management of aquaculture farming is of great concern due to possible residues in aquatic products and in the development of antibiotic resistance. In 2005 the Vietnamese government first promulgated a list of medicines, chemicals and antibiotics that are banned or limited for use in aquaculture and this is regularly updated and amended to tackle the growing problem of antibiotic resistance.

The fishery sector in Vietnam

Fisheries is a key national economic sector that produces 4–5% GDP and 5–6% of the total national turnover of Vietnam. Aquaculture in Vietnam has changed from a self-sufficient sector in the early 1960s

into a concentrated commodity production and this model has become the most important contributor in the fishery sector since 2007. In Vietnam, the Mekong River Delta in the south and the Red River Delta in the north have been the major areas for wild caught fishing as well as intensive aquaculture farming for decades. Aquaculture production in the Mekong Delta has contributed more than 70% of the country's total production, in which catfish (fresh water) and black tiger and white leg shrimp (brackish water) are the three most important commercial species grown (Figures 1, 2). Since the 1990s, fishery products have been the third biggest export commodity of the country. Fishery exports earned over US\$1 billion in 2000, US\$3 billion in 2006, US\$6 billion in 2011 and reached a peak of US\$7.9 billion in 2014. Currently, Vietnamese fishery products are exported to more than 140 countries and territories in five continents. According to the Food and Agriculture Organisation of the United Nations (FAO), nearly 10% of the population in Vietnam obtains their main income from Fisheries. More than four million workers including a large proportion of women are employed in aquaculture and fishery activities (Figure 3).

Antibiotic resistance in aquaculture in Vietnam

In Vietnam, only limited studies on antibiotic uses and resistance in aquaculture were conducted and/or published before 2005. A study on antibiotic residues and resistance against Norfloxacin (NOR), Oxolinic acid (OXO), Trimethoprim (TRI) and Sulfamethoxazole



Figure 1. Seabass or Barramundi (*Lates calcarifer*) farming in Vietnam.



Figure 2. Tra catfish feeding in a farm in the Mekong Delta, Vietnam.

(SUL) has been conducted in four shrimp farming locations (Thai Binh, Nam Dinh, Can Gio-Ho Chi Minh City and Ca Mau provinces) in mangrove areas in Vietnam in 2003¹. The results showed the presence of antibiotic residues and a high incidence of bacterial resistance to the four antibiotics observed in these shrimp ponds in different mangrove areas in Vietnam. Another study on antibiotic resistance of 123 bacterial isolates (from water, sediment and different fish farms (catfish, tilapia, common carp and gourami) in five provinces in the Mekong Delta) found that 90% of the isolates were resistant to Tetracycline (TET), 76% to Ampicillin (AMP), 100% to Chloramphenicol (CHL), 65% to Nitrofurantoin (NIT) and 89% to Trimethoprim-Sulphamethoxazole (SXT).

A number of extensive research projects on antibiotic resistance of commensal bacteria from several aquaculture species have been conducted by our groups. In 2003, a study on the resistance of 101 isolates (belonging to the family Enterobacteriaceae) from frozen catfish of the Mekong Delta showed high frequencies of resistance (80–100%) to Oxytetracycline (OTE), CHL, Streptomycin (STR), SXT and AMP (unpublished data). The levels of multiple resistance were also high, of which 100% of isolates were resistant to at least four antibiotics. A research study published in 2005 reported the resistance of 92 isolates (including Enterobacteriaceae (49.1%), *Pseudomonas* spp. (35.2%) and *Vibrionaceae* (15.7%) families) from three catfish farms in the Mekong Delta². Rather high incidences



Figure 3. Tra catfish filleting in a seafood processing company in the Mekong Delta, Vietnam.

(20–60%) of resistance to OTE, CHL, SXT, NIT, Nalidixic acid (NAL) and AMP were observed. The multiple antibiotic resistance (MAR) index values indicated that the three farms corresponded to high-risk exposed-antibiotic sources. In a recent publication, a larger collection of samples (catfish, water and sediment) was collected from 50 catfish culture ponds (belonging to 15 catfish intensive farms) in the Mekong Delta, Vietnam³. All 116 *Pseudomonas* spp. and 92 *Aeromonas* spp. isolates recovered from these samples were tested for their resistance to 13 antimicrobial agents belonging to eight families. High levels of resistance to AMP, SXT, and NAL were observed. Lower frequencies of resistance occurred against Ciprofloxacin (CIP), NOR, TET, Doxycycline (DOX), Gentamycin (GEN), Neomycin (NEO), STR, and Kanamycin (KAN). The percentages of multiple resistance (resistance to three or more antimicrobial classes) was 96.6% for *Pseudomonas* and 61.9% for *Aeromonas* spp. The MAR indices showed that *Pseudomonas* and *Aeromonas* isolates were high-risk sources of contamination where antibiotics were commonly used. The level of resistance to CHL and NIT was still high despite these antibiotics being banned for use in aquaculture in Vietnam since 2005, and this resistance is possibly due to the co-transfer of resistance genes. Our recent study on the resistance of 167 *Vibrio* spp. isolated from blood cockles (16 farms of the six provinces in the Mekong Delta) showed a low incidence of resistance to nine tested antibiotics (TET, NAL, NOR, CIP, SXT, NIT, CHL, STR, KAN), with a high level of resistance to AMP only. Unfortunately, very high levels of resistance of bacteria in aquaculture environments or products have also been reported in many countries which have intensive aquaculture activities such as China⁴, USA⁵, Thailand⁶, India⁷, Indonesia⁸, Australia⁹, Italy¹⁰ and Chile¹¹ implying the heavy use of antibiotics in aquaculture worldwide and this provides a warning on the effects of antibiotic usage in aquatic ecosystems, and the world wide emergence of antibiotic resistance in aquatic bacteria.

Integrations and the transferability of antibiotic resistant genes

In bacteria, horizontal gene transfer is the principle mechanism responsible for the spread of antibiotic resistance genes¹². Horizontal gene transfer is facilitated by mobile genetic elements such as conjugative plasmids, transposons, and phages¹³. Integrations are not themselves mobile elements but are associated with mobile genetic elements (transposons or conjugative plasmids) enabling efficient intra- or interspecies transmission¹⁴. In Vietnam, studies on integrations and resistance gene transfer mechanisms have been conducted in clinical isolates^{1,15–19}. However, very limited studies have been published on non-human isolates such as food-borne pathogens²⁰, animal pathogens¹⁹ and catfish pathogens²¹. Recently we have done an extensive study on integrations and the transferability of antibiotic resistant genes of commensal bacteria in catfish aquaculture in Vietnam³. We found that the commensal isolates of catfish harbour a pool of mobile genetic elements such as plasmids and integrations, which contain various antibiotic resistance gene cassettes. Conjugation and transformation experiments demonstrated the successful transfer of all or part of the resistance phenotypes of isolates to the recipient strains of different genera and sources. Class 1 integrations associated with plasmids have facilitated the emergence and dissemination of antibiotic resistance in aquaculture environments. The transformation and conjugation experiments also indicated that CHL resistance phenotype was co-transferred in association with SXT, AMP, TET resistance phenotypes. In addition, the CHL resistance genes (*catB*) have often been associated with class 1 integrations, therefore, the co-selection is important for CHL resistance dissemination. This suggests that, in the absence of chloramphenicol selection pressure, CHL resistance is co-transferred and maintained due to gene linkage to genes encoding resistance to antimicrobials that are widely used in food animals or aquaculture²². The other studies worldwide also indicate that aquatic environments or products are reservoirs of antibiotic resistance and class 1 integrations have facilitated the dissemination of antibiotic resistant genes such as in the USA²³, China²⁴, Thailand²⁵, Northern Europe and North America²⁶, Europe, Japan and the USA²⁷, and Japan²⁸.

Conclusion

The high frequencies of resistance observed in aquaculture bacterial isolates in Vietnam, and in worldwide aquatic environmental and product isolates, reflect the global spread of resistance due to extensive use of antibiotics in aquaculture. Aquaculture resident bacteria could serve as a reservoir of resistant genes if they harbour a pool of mobile genetic elements that can readily be transferred intra- and interspecies.

References

1. Le, T.X. *et al.* (2005) Antibiotic resistance in bacteria from shrimp farming in mangrove areas. *Sci. Total Environ.* **349**, 95–105. doi:10.1016/j.scitotenv.2005.01.006
2. Sarter, S. *et al.* (2007) Antibiotic resistance in Gram-negative bacteria isolated from farmed catfish. *Food Contr.* **18**, 1391–1396. doi:10.1016/j.foodcont.2006.10.003
3. Nguyen, H.N.K. *et al.* (2014) Molecular characterization of antibiotic resistance in *Pseudomonas* and *Aeromonas* isolates from catfish of the Mekong Delta, Vietnam. *Vet. Microbiol.* **171**, 397–405. doi:10.1016/j.vetmic.2014.01.028
4. Su, H.-C. *et al.* (2011) Occurrence of antibiotic resistance and characterization of resistance genes and integrons in *Enterobacteriaceae* isolated from integrated fish farms in south China. *J. Environ. Monit.* **13**, 3229–3236. doi:10.1039/c1em10634a
5. McPhearson, R.M. *et al.* (1991) Antibiotic resistance in Gram-negative bacteria from cultured catfish and aquaculture ponds. *Aquaculture* **99**, 203–211. doi:10.1016/0044-8486(91)90241-X
6. Kitiyodom, S. *et al.* (2010) Characterization of antibiotic resistance in *Vibrio* spp. isolated from farmed marine shrimps (*Penaeus monodon*). *FEMS Microbiol. Ecol.* **72**, 219–227. doi:10.1111/j.1574-6941.2010.00846.x
7. Vivekanandhan, G. *et al.* (2002) Antibiotic resistance of *Aeromonas hydrophila* isolated from marketed fish and prawn of South India. *Int. J. Food Microbiol.* **76**, 165–168. doi:10.1016/S0168-1605(02)00009-0
8. Yusop, Z. *et al.* (2009) Antibiotic resistance and plasmid profiling of *Vibrio parahaemolyticus* isolated from cockles in Padang, Indonesia. *Int. Food. Res. J.* **16**, 53–58.
9. Akinbowale, O.L. *et al.* (2006) Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. *J. Appl. Microbiol.* **100**, 1103–1113. doi:10.1111/j.1365-2672.2006.02812.x
10. Ottaviani, D. *et al.* (2001) Antimicrobial susceptibility of potentially pathogenic halophilic vibrios isolated from seafood. *Int. J. Antimicrob. Agents* **18**, 135–140. doi:10.1016/S0924-8579(01)00358-2
11. Miranda, C.D. and Zemelman, R. (2002) Antimicrobial multiresistance in bacteria isolated from freshwater Chilean salmon farms. *Sci. Total Environ.* **293**, 207–218. doi:10.1016/S0048-9697(02)00022-0
12. de la Cruz, F. and Davies, J. (2000) Horizontal gene transfer and the origin of species: lessons from bacteria. *Trends Microbiol.* **8**, 128–133. doi:10.1016/S0966-842X(00)01703-0
13. Smets, B.F. and Barkay, T. (2005) Horizontal gene transfer: perspectives at a crossroads of scientific disciplines. *Nat. Rev. Microbiol.* **3**, 675–678. doi:10.1038/nrmicro1253
14. Davies, J. and Davies, D. (2010) Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol. Rev.* **74**, 417–433. doi:10.1128/MMBR.00016-10
15. Dalsgaard, A. *et al.* (1999) Cholera in Vietnam: changes in genotypes and emergence of class I integrons containing aminoglycoside resistance gene cassettes in *Vibrio cholerae* O1 strains isolated from 1979 to 1996. *J. Clin. Microbiol.* **37**, 734–741.
16. Iversen, J. *et al.* (2003) Characterization of antimicrobial resistance, plasmids, and gene cassettes in *Sbigella* spp. from patients in Vietnam. *Microb. Drug Resist.* **9**, 17–24. doi:10.1089/107662903322541856
17. Le, V. *et al.* (2015) Genetic characterization of three *qnrSI*-harbouring multidrug-resistance plasmids and *qnrSI*-containing transposons circulating in Ho Chi Minh City, Vietnam. *J. Med. Microbiol.* **64**, 869–878. doi:10.1099/jmm.0.000100
18. Naas, T. *et al.* (2000) Integron-located VEB-1 extended-spectrum β -lactamase gene in a *Proteus mirabilis* clinical isolate from Vietnam. *J. Antimicrob. Chemother.* **46**, 703–711. doi:10.1093/jac/46.5.703
19. Vo, A.T.T. *et al.* (2010) Antimicrobial resistance, class I integrons, and genomic Island 1 in *Salmonella* isolates from Vietnam. *PLoS One* **5**, e9440. doi:10.1371/journal.pone.0009440
20. Van, T.T.H. *et al.* (2007) Antibiotic resistance in food-borne bacterial contaminants in Vietnam. *Appl. Environ. Microbiol.* **73**, 7906–7911. doi:10.1128/AEM.00973-07
21. Dung, T.T. *et al.* (2009) IncK plasmid-mediated tetracycline resistance in *Edwardsiella ictaluri* isolates from diseased freshwater catfish in Vietnam. *Aquaculture* **295**, 157–159. doi:10.1016/j.aquaculture.2009.07.010
22. Bischoff, K.M. *et al.* (2005) The chloramphenicol resistance gene *cmlA* is disseminated on transferable plasmids that confer multiple-drug resistance in swine *Escherichia coli*. *FEMS Microbiol. Lett.* **243**, 285–291. doi:10.1016/j.femsle.2004.12.017
23. Nawaz, M. *et al.* (2010) Detection and characterization of virulence genes and integrons in *Aeromonas veronii* isolated from catfish. *Food Microbiol.* **27**, 327–331. doi:10.1016/j.fm.2009.11.007
24. Chen, B. *et al.* (2011) Class I integrons, selected virulence genes, and antibiotic resistance in *Escherichia coli* isolates from the Minjiang River, Fujian Province, China. *Appl. Environ. Microbiol.* **77**, 148–155. doi:10.1128/AEM.01676-10
25. Agersø, Y. and Petersen, A. (2007) The tetracycline resistance determinant Tet 39 and the sulphonamide resistance gene *sullI* are common among resistant *Acinetobacter* spp. isolated from integrated fish farms in Thailand. *J. Antimicrob. Chemother.* **59**, 23–27. doi:10.1093/jac/dkl419
26. Schmidt, A.S. *et al.* (2001) Characterization of class I integrons associated with R-plasmids in clinical *Aeromonas salmonicida* isolates from various geographical areas. *J. Antimicrob. Chemother.* **47**, 735–743. doi:10.1093/jac/47.6.735
27. L'Abée-Lund, T.M. and Sørum, H. (2001) Class I integrons mediate antibiotic resistance in the fish pathogen *Aeromonas salmonicida* worldwide. *Microb. Drug Resist.* **7**, 263–272. doi:10.1089/10766290152652819
28. Ahmed, A.M. *et al.* (2015) Seafood as a reservoir of Gram-negative bacteria carrying integrons and antimicrobial resistance genes in Japan. *Biomed. Environ. Sci.* **28**, 924–926.

Biographies

Dr Hoang Nam Kha Nguyen is a lecturer in Department of Aquatic Product Processing at the Faculty of Fisheries at Nong Lam University. His research interests are aquaculture commensal bacteria and molecular biology.

Dr Thi Thu Hao Van is a Research Fellow in the School of Science, RMIT University. Her research interests are antibiotic resistance and virulence of bacterial pathogens, the development of bacteria from gut microbiota for use as probiotics, identifying the cause of disease in farm animals, and the development of vaccines against the pathogens causing these diseases.

Professor Peter J Coloe is Pro Vice-Chancellor, College of Science, Engineering and Health, RMIT University and is a Fellow of ASM. His research interests are in vaccines and disease diagnostics and novel ways to control disease and to deliver vaccine antigens.